



Politecnico di Torino

Porto Institutional Repository

[Article] Killing vector fields of constant length on compact hypersurfaces

Original Citation:

Di Scala A.J. (2015). *Killing vector fields of constant length on compact hypersurfaces*. In: [GEOMETRIAE DEDICATA](#), vol. 175 n. 1, pp. 403-406. - ISSN 0046-5755

Availability:

This version is available at : <http://porto.polito.it/2535724/> since: March 2014

Publisher:

Springer

Published version:


DOI:[10.1007/s10711-014-9954-6](https://doi.org/10.1007/s10711-014-9954-6)

Terms of use:

This article is made available under terms and conditions applicable to Open Access Policy Article ("Public - All rights reserved") , as described at http://porto.polito.it/terms_and_conditions.html

Porto, the institutional repository of the Politecnico di Torino, is provided by the University Library and the IT-Services. The aim is to enable open access to all the world. Please [share with us](#) how this access benefits you. Your story matters.

(Article begins on next page)



KILLING VECTOR FIELDS OF CONSTANT LENGTH ON COMPACT HYPERSURFACES

ANTONIO J. DI SCALA

Abstract: We show that if a compact hypersurface $M \subset \mathbb{R}^{n+1}$, $n \geq 3$, admits a non zero Killing vector field X of constant length then n is odd and M is diffeomorphic to the unit hypersphere of \mathbb{R}^{n+1} . Actually, we show that M is a complex ellipsoid in $\mathbb{C}^N = \mathbb{R}^{n+1}$. As an application we give a simpler proof of a recent theorem due to S. Deshmukh [2].

1. INTRODUCTION

In a recent paper [2], S. Deshmukh motivated by the Hopf vector fields of odd dimensional hyperspheres asked the following question:

“Does an orientable compact hypersurface in a Euclidean space that admits a unit Killing vector field ξ together with the condition that $g(A\xi, \xi)$ is a constant necessarily isometric to a sphere?^[sic]”

Deshmukh answer positively but his proof is not correct. In [2, page 972] after equation (7) he claims that the shape operator A of a hypersurface M satisfies

$$d\psi_t \circ A = A \circ d\psi_t$$

where $\{\psi_t\}$ is the flow of the unit Killing vector field ξ . This claim is false since the shape operator A contains extrinsic information about M . For example, if M is a plane curve and t its arc length parameter then $\xi = \partial_t$ is a unit Killing vector field. But then the above equation implies that the curvature (i.e. the inverse of the radius of the osculating circle) of the plane curve M is constant which is not true for all plane curves.

It is known that compact hypersurfaces are orientable (see for example [5]) so such hypothesis can be omitted from the above question.

In this article we show that besides Deskmush’s wrong proof the answer to his question is indeed positive. Moreover, we show that the condition $g(A\xi, \xi)$ to be constant, where g is the induced Riemannian

Date: March 23, 2015.

AMS CLASSIFICATION: 53C20, 53C22, 53C25.

metric on M , is not necessary to characterize M topologically and geometrically.

Here is our main result.

Theorem 1.1. *A compact hypersurface $M \subset \mathbb{R}^{n+1}$ that admits a unit Killing vector field ξ is diffeomorphic to an odd dimensional hypersphere. Actually, $n + 1 = 2N$ and M is (up to isometry $\mathbb{R}^{n+1} \cong \mathbb{C}^N$) the complex ellipsoid defined by the equation*

$$|\omega_1 z_1|^2 + |\omega_2 z_2|^2 + \cdots + |\omega_N z_N|^2 = 1$$

where (z_1, \dots, z_N) are complex coordinates of \mathbb{C}^N and $\omega_1, \dots, \omega_N$ are non zero real numbers. Moreover, if $g(A\xi, \xi)$ is constant then

$$\omega_1^2 = \cdots = \omega_N^2$$

and M is a hypersphere of constant curvature.

For more about Killing vector fields of constant length on Riemannian manifolds see [1].

2. PRELIMINARIES

A Killing vector field X of the flat Euclidean space \mathbb{R}^{n+1} is given by a pair $X = (R, \mathbf{v})$ where R is a skew-symmetric matrix and $\mathbf{v} \in \mathbb{R}^{n+1}$ is a vector. At a point $\mathbf{p} \in \mathbb{R}^{n+1}$, $X(\mathbf{p})$ is given by the formula

$$X(\mathbf{p}) = R\mathbf{p} + \mathbf{v}.$$

The block diagonal form for skew-symmetric matrices can be used to introduce complex coordinates (z_1, \dots, z_N) on the image of R . So, we have that $\mathbb{R}^{n+1} = \mathbb{C}^N \oplus \mathbb{R}^k$ and the Killing vector field X is given by

$$(1) \quad X(z_1, \dots, z_N, \mathbf{r}) = (i\omega_1 z_1, \dots, i\omega_N z_N, \mathbf{v}_k)$$

where \mathbf{v}_k is the projection of \mathbf{v} to \mathbb{R}^k and $\omega_1, \dots, \omega_N$ are non zero real numbers.

Let $H_X^\mathcal{L} \subset \mathbb{R}^{n+1}$, $\mathcal{L} > 0$, be the set of points where the Killing vector field X has constant length \mathcal{L} . The following is a straightforward consequence of equation (1).

Proposition 2.1. *The set $H_X^\mathcal{L} \subset \mathbb{R}^{n+1}$ is a smooth connected submanifold and the Killing vector field X is always tangent to $H_X^\mathcal{L}$. Moreover, if $H_X^\mathcal{L} \neq \emptyset$ then one of the following holds:*

- (i) $\mathcal{L} = \|\mathbf{v}_{ker}\|$ and $H_X^\mathcal{L} = \mathbb{R}^k$,
- (ii) $\mathcal{L} > \|\mathbf{v}_{ker}\|$ and $H_X^\mathcal{L} = \mathbf{E} \times \mathbb{R}^k \subset \mathbb{C}^N \times \mathbb{R}^k$ where $\mathbf{E} \subset \mathbb{C}^N$ is the complex ellipsoid defined by

$$|\omega_1 z_1|^2 + \cdots + |\omega_N z_N|^2 = \mathcal{L} - \|\mathbf{v}_{ker}\|,$$

where $\omega_1, \dots, \omega_N$ are non zero real numbers.

Here is another important consequence of equation (1).

Lemma 2.2. *The rank of the shape operator S_p of the hypersurface $H_X^\mathcal{L}$ is constant i.e. it does not depend upon the point $p \in H_X^\mathcal{L}$. Moreover, in case (ii) of the previous proposition the rank of S is $2N - 1$ at all points of $H_X^\mathcal{L}$.*

Proof. Indeed, the vector field $\mathbf{n} = (\omega_1^2 z_1, \dots, \omega_N^2 z_N)$ is perpendicular to the complex ellipsoid E at $p = (z_1, \dots, z_N)$. Then for $\mathbf{t} \in T_p E$ we have

$$S_p(\mathbf{t}) \cdot \mathbf{t} = -D_{\mathbf{t}} \mathbf{n} \cdot \mathbf{t} = - \sum_{j=1}^N \omega_j^2 |t_j|^2.$$

Since $\omega_1, \dots, \omega_N$ are non zero real numbers we get that the rank of the shape operator S_p is $2N - 1$. \square

3. MAIN THEOREM

Theorem 3.1. *Let (M^n, g) be a connected compact Riemannian manifold of dimension $n \geq 3$. Let ξ be a non zero Killing vector field of M with constant length $\mathcal{L} = \|\xi\|$.*

If (M, g) can be isometrically embedded into the flat Euclidean space \mathbb{R}^{n+1} then n is odd and

- (i) *the Killing vector field ξ extends to a Killing vector field $X = (R, \mathbf{v})$ of \mathbb{R}^{n+1} ,*
- (ii) *$M = H_X^\mathcal{L}$.*

Proof. We will identify M with its image under the isometric embedding into \mathbb{R}^{n+1} . The idea is that for $n \geq 3$ hypersurfaces of \mathbb{R}^{n+1} whose shape operator has rank ≥ 3 are rigid due to the theorem of Beez-Killing [4, page 46, Corollary 6.5].

Let A be the shape operator of M . It is well known that M has a point q where A_q is positive (or negative) definite [3, page 255]. Since the rank of A_q is n the theorem of Beez-Killing implies that the Killing vector field ξ extends to a Killing vector field X of \mathbb{R}^{n+1} near q . Then near q the hypersurface M is contained in $H_X^\mathcal{L}$. If $\dim(H_X^\mathcal{L}) = n + 1$ then X is a parallel vector field i.e. $X = \mathbf{v}$ a constant vector. Then ξ is the restriction of a constant vector and so $A_q(\xi) = 0$ which contradicts that A_q is definite at q . So $\dim(H_X^\mathcal{L}) = n$ and $M = H_X^\mathcal{L}$ near q . Notice that $H_X^\mathcal{L}$ is as in (ii) of Proposition 2.1. That is to say, the shape operator of $H_X^\mathcal{L}$ has rank $n = 2N - 1$ at q which show that n is odd.

Let $U = M \cap H_X^\mathcal{L}$ be the intersection of both hypersurfaces and let $\overset{\circ}{U}$ be the interior of U in M . By the previous observation $\overset{\circ}{U}$ contains a neighborhood of q in M and so $\overset{\circ}{U}$ is non empty. We claim that $\overset{\circ}{U}$ is closed in M . Indeed, let $p_j \in \overset{\circ}{U}$ be a sequence converging to $p_0 \in M$. The

shape operator A_{p_0} is the limit of the shape operator S_{p_j} of $H_X^{\mathcal{L}}$. Then $A_{p_0} = S_{p_0}$ and Lemma 2.2 implies that the rank of A_{p_0} is $n \geq 3$. Again by the theorem of Beez-Killing we have that near p_0 the vector field ξ extends to a Killing vector field Y of \mathbb{R}^{n+1} . Notice that Y restricted to $\overset{\circ}{U}$ agree with X near the points p_j for $j \gg 0$. This implies $X = Y$ on \mathbb{R}^{n+1} and then $p_0 \in \overset{\circ}{U}$. So $\overset{\circ}{U} = M$ hence $M \subset H_X^{\mathcal{L}}$. This shows (i) and (ii) since by Proposition 2.1 the compact hypersurface $H_X^{\mathcal{L}}$ is connected. \square

3.1. Proof of Theorem 1.1. It is clear that the first part of Theorem 1.1 follows from Theorem 3.1. By using the shape operator as in the proof of Lemma 2.2 the condition $g(A\xi, \xi)$ is constant gives:

$$\omega_1^4 |z_1|^2 + \cdots + \omega_N^4 |z_N|^2 = cte,$$

on the complex ellipsoid $|\omega_1 z_1|^2 + \cdots + |\omega_N z_N|^2 = 1$. Then

$$\omega_1^4 |z_1|^2 + \cdots + \omega_N^4 |z_N|^2 = cte (|\omega_1 z_1|^2 + \cdots + |\omega_N z_N|^2)$$

for all $(z_1, \dots, z_N) \in \mathbb{C}^N$. So $\omega_j^2 = cte$ for $j = 1, \dots, N$ and M is the hypersphere of \mathbb{C}^N defined by the equation

$$|z_1|^2 + \cdots + |z_N|^2 = \frac{1}{cte}.$$

\square

4. FURTHER REMARKS

Surfaces in \mathbb{R}^3 are not rigid so it is natural to ask about existence of compact surfaces with a unit Killing vector field in \mathbb{R}^3 . The answer is that such surface does not exist. Indeed, it is not difficult to see that a unit Killing vector field of a surface is a parallel vector field. So the metric is flat. But a flat compact surface cannot be a surface in \mathbb{R}^3 due to the classical theorem about existence of elliptic points [3, page 255].

REFERENCES

- [1] BERESTOVSKII, V.N. AND NIKONOROV, YU. G.: Killing vector fields of constant length on Riemannian manifolds. *Sibirsk. Mat. Zh.* 49 (2008), no. 3, 497–514; translation in *Sib. Math. J.* 49 (2008), no. 3, 395–407 <http://rd.springer.com/article/10.1007%2Fs11202-008-0039-3>
- [2] DESHMUKH, S.: A note on compact hypersurfaces in a Euclidean space. *C. R. Math. Acad. Sci. Paris* 350 (2012), no. 21–22, 971–974. <http://www.sciencedirect.com/science/article/pii/S1631073X12003172>
- [3] GALLOT, S. ; HULIN, D. AND LAFONTAINE, J.: Riemannian Geometry, third Edition, Springer (2004).
- [4] KOBAYASHI, S. AND NOMIZU, K.: Foundations of differential geometry. Vol II. Interscience Publishers, a division of John Wiley & Sons, New York-London (1969).
- [5] SAMELSON, H.: Orientability of hypersurfaces in \mathbb{R}^n . *Proc. Amer. Math. Soc.* 22, 1969, 301–302. <http://www.ams.org/journals/proc/1969-022-01/S0002-9939-1969-0245026-9/>

Antonio J. Di Scala,
Dipartimento di Scienze Matematiche “G. L. Lagrange”,
Politecnico di Torino, Corso Duca degli Abruzzi 24,
10129 Torino, Italy.
email: antonio.discal@polito.it