



# Politecnico di Torino

## Porto Institutional Repository

[Article] Tsallis and Kaniadakis Entropic Measures in Stellar Polytropes

*Original Citation:*

A.C. Sparavigna (2015). *Tsallis and Kaniadakis Entropic Measures in Stellar Polytropes*. In: [PHILICA](#), vol. 2015 n. 541, pp. 1-4. - ISSN 1751-3030

*Availability:*

This version is available at : <http://porto.polito.it/2621751/> since: November 2015

*Publisher:*

PHILICA

*Terms of use:*

This article is made available under terms and conditions applicable to Open Access Policy Article ("Creative Commons: Attribution 3.0") , as described at [http://porto.polito.it/terms\\_and\\_conditions.html](http://porto.polito.it/terms_and_conditions.html)

Porto, the institutional repository of the Politecnico di Torino, is provided by the University Library and the IT-Services. The aim is to enable open access to all the world. Please [share with us](#) how this access benefits you. Your story matters.

*Publisher copyright claim:*

This is the publisher version of an article published on [PHILICA](#), Publisher [pin missing: publisher], Vol 2015 , Number 541 Year 2015 (ISSN [pin missing: issn] )The present version is accessible on PORTO, the Open Access Repository of the Politecnico of Torino

(Article begins on next page)

[Philica front page](#)[Search](#)[About Philica](#)[Take the tour](#)[Publish your work](#)[Work needing review](#)[Most popular entries](#)[Highest-rated entries](#)[Recent reviews](#)[How to cite Philica](#)[FAQs](#)[Support Philica](#)[Contact us](#)[Get confirmed status](#)

## Tsallis and Kaniadakis Entropic Measures in Stellar Polytropes

[Amelia Carolina Sparavigna](#) (Department of Applied Science and Technology, Politecnico di Torino)

Published in [astro.philica.com](http://astro.philica.com)

### Abstract

Polytropes are self-gravitating fluid spheres used in astrophysics as crude approximation of more realistic stellar models. They possess equations that have scale parameters linked to mass, energy and entropy. Since Boltzmann distribution yields unphysical results, the use of generalized entropies, such as Tsallis and Kaniadakis entropies, had been proposed. Here we discuss how these entropies are related in polytrope solutions.

### Article body

## Tsallis and Kaniadakis Entropic Measures in Stellar Polytropes

**Amelia Carolina Sparavigna**

Department of Applied Science and Technology, Politecnico di Torino, Torino, Italy

**Abstract:** Polytropes are self-gravitating fluid spheres used in astrophysics as crude approximation of more realistic stellar models. They possess equations that have scale parameters linked to mass, energy and entropy. Since Boltzmann distribution yields unphysical results, the use of generalized entropies, such as Tsallis and Kaniadakis entropies, had been proposed. Here we discuss how these entropies are related in polytrope solutions.

**Keywords:** Entropy, Generalized Entropies.

### 1. Introduction

In astrophysics, a polytrope refers to a solution of the Lane–Emden equation. This is an equation which gives the pressure as a function of density [1]. These solutions are modelling self-gravitating fluid spheres that are called “polytropes” too, which are objects used as crude approximation to more realistic stellar models [2]. The solution of the Lane–Emden equation, a dimensionless form of Poisson's equation for the gravitational potential, depends on a parameter which is the polytropic index  $n$ . It is written as  $P=K\rho^{(n+1)/n}$ , where  $P$  is pressure,  $\rho$  is density and  $K$  a constant. If stellar structure is approximated with a polytrope having a given index, then two scaling parameters are needed to express the structure in physical units [3]. The two parameters that we can use are a constant which is related to entropy and the stellar mass. Since Boltzmann distribution yields unphysical results, the Boltzmann entropy had been substituted by a generalized entropy, the Tsallis entropy [4]. Another generalized entropy, the Kaniadakis entropy, had been recently proposed too, in [5]. Here we discuss how these two entropies are related in polytrope solutions, and that the result given in [5] can be easily obtained from [4].

### 2. The entropies

Well-known is the entropy proposed by Claude Shannon in 1948 [6]. He defined the entropy  $H$  of a discrete random variable  $X$ , as the expected value of the information content:  $H(X)=-\sum_i p_i \log_b p_i$ . The probability of  $i$ -event is  $p_i$  and  $b$  is the base of the used logarithm. However, several entropies exist which are generalizing Shannon entropy. Among them we have Tsallis and Kaniadakis entropies [7,8], which are defined, with a corresponding choice of

NEWS: The SOAP Project, in collaboration with CERN, are conducting a [survey on open-access publishing](#). Please take a moment to give them your views

measurement units equal to 1, as follow:

$$(1) \quad T_{\text{Tsallis}} : T = T_q = \frac{1}{q-1} \left( 1 - \sum_i p_i^q \right)$$

$$(2) \quad \text{Kaniadakis } (\kappa\text{-entropy}) : K_\kappa = - \sum_i \frac{p_i^{1+\kappa} - p_i^{1-\kappa}}{2\kappa}$$

In (1) and (2) we have the entropic indices  $q$  and  $\kappa$ . For its generalized additivity, the Kaniadakis entropy requires another function, defined as follow:

$$\mathfrak{S} = \sum_i (p_i^{1+\kappa} + p_i^{1-\kappa}) / 2$$

A detailed discussion of the generalized additivity of Tsallis and  $\kappa$ -entropy is given in [9].

Tsallis and Kaniadakis entropies are linked:

$$(3) \quad K_\kappa = \frac{T_{1+\kappa} + T_{1-\kappa}}{2},$$

$$\text{where } T(q=1+\kappa) = -\frac{1}{\kappa} \sum_i p_i^{1+\kappa} + \frac{1}{\kappa}; \quad T(q=1-\kappa) = \frac{1}{\kappa} \sum_i p_i^{1-\kappa} - \frac{1}{\kappa}$$

Eq.(3) is a simpler form of an expression given in [10,11]. However, besides this relation, because of the generalized additivity possessed by the Kaniadakis entropy, we need also another relation:

$$(4) \quad \mathfrak{S}_\kappa = \frac{\kappa}{2} \left( -T_{1+\kappa} + T_{1-\kappa} + \frac{2}{\kappa} \right)$$

In (3) and (4), we have Kaniadakis functions expressed by Tsallis entropy. As shown in [12], we can also write  $T$  expressed by means of Kaniadakis functions:

$$2K + \frac{2}{\kappa} \mathfrak{S} = T_{1+\kappa} + T_{1-\kappa} + \left( -T_{1+\kappa} + T_{1-\kappa} + \frac{2}{\kappa} \right)$$

$$\text{And then : (5) } K_\kappa + \frac{1}{\kappa} \mathfrak{S}_\kappa = T_{1-\kappa} + \frac{1}{\kappa}$$

Let us have:  $\kappa=1-q$ . From (5) we have immediately the relation between Tsallis and Kaniadakis functions:

$$(6) \quad T_q = K_{1-q} + \frac{\mathfrak{S}_{1-q} - 1}{(1-q)}$$

### 3. With polytropes

The relation (6) between Tsallis and Kaniadakis entropies can be useful in several problems. Here we consider its use in polytropes. In the previous equations, we have  $p_i$  denoting the probability distribution. In Ref.4, it is used letter  $f$  for probability. From now on, we will use this notation. In [4], the distribution from Tsallis entropy is:

$$(7) \quad C_q(f) = \frac{f(1-f^{q-1})}{q-1}$$

After Eq.6, we can write Eq.7 in the following manner:

We aim to suit all browsers, but recommend Firefox particularly:



$$(8) \quad C_q(f) = \frac{f - f^q}{q-1} = \frac{1}{2(q-1)}(f^{2-q} - f^q) - \frac{1}{2(q-1)}(f^{2-q} + f^q) + \frac{f}{q-1}$$

Of course, (7) and (8) are the same equation. As a consequence, Kaniadakis distributions are linked to Tsallis distribution by:

$$(9) \quad t_q(f) = C_q(f) = \frac{(f - f^q)}{q-1} = k_{1-q} - g_{1-q} + \frac{f}{q-1}$$

where  $K_{1-q} = \sum_i k_{1-q}$ ,  $\mathfrak{S}_{1-q} = \sum_i g_{1-q}$ ,  $1 = \sum_i f$

From [4], a relation exists between polytrope index and entropic Tsallis index:

$$(10) \quad n = \frac{3}{2} + \frac{1}{q-1}$$

As a special case, for  $q \rightarrow 1$ , we find the isothermal situation. To have Eq.6, as shown in [12], we need  $\kappa=1-q$  or  $\kappa=q-1$ . Then, from (10), considering that we have for the Kaniadakis index,  $-1 < \kappa < 1$ :

$$(11) \quad n = \frac{3}{2} + \frac{1}{\kappa}$$

And in fact, (11) is the relation that we find in [5]. Using then the relation between Tsallis and Kaniadakis entropies and distributions we can easily find results concerning several applications. Polytropes are an example of such a possible approach.

## References

- [1] Horedt, G.P. (2004). Polytropes. Applications in Astrophysics and Related Fields, Kluwer. ISBN 1-4020-2350-2
- [2] Chandrasekhar, S. (1958). An Introduction to the Study of Stellar Structure, Dover.
- [3] Knapp, J. (2015). Polytropes, retrieved 9 November 2015 from <http://www.astro.princeton.edu/~gk/A403/polytrop.pdf>
- [4] Plastino, A.R.; Plastino, A. (1993). Stellar Polytropes and Tsallis Entropy. Physics Letters A 174(5):384-386. DOI: 10.1016/0375-9601(93)90195-6
- [5] Bento, E.P.; Silva, J.R.P.; Silva, R. (2013). Non-Gaussian Statistics, Maxwellian Derivation and Stellar Polytropes. Physica A: Statistical Mechanics and its Applications 392.4 (2013): 666-672. DOI: 10.1016/j.physa.2012.10.022
- [6] Shannon, C.E. (1948). A Mathematical Theory of Communication. Bell System Technical Journal 2(3):379-423. DOI: 10.1002/j.1538-7305.1948.tb01338.x
- [7] Tsallis, C. (1960). Possible Generalization of Boltzmann-Gibbs Statistics, Journal of Statistical Physics, 1988, 52: 479-487. DOI:10.1007/BF01016429
- [8] Kaniadakis, G.(2002). Statistical Mechanics in the Context of Special Relativity, Phys. Rev. E, 2002, 66, 056125. DOI: 10.1103/physreve.66.056125
- [9] Sparavigna, A.C. (2015). On the Generalized Additivity of Kaniadakis Entropy, Int. J. Sci. 4(2):44-48. DOI: 10.18483/ijSci.627
- [10] Kaniadakis, G. (2001). Non-Linear Kinetics Underlying Generalized Statistics, Physica A 296(3-4):405-425. DOI: 10.1016/s0378-4371(01)00184-4
- [11] Santos, P.; Silva R.; Alcaniz J.S.; Anselmo, D.H.A.L. (2011). Generalized Quantum

Entropies, Physics Letters A 375:3119–3123. DOI: 10.1016/j.physleta.2011.07.001

[12] Sparavigna, A.C. (2015). Relations Between Tsallis and Kaniadakis Entropic Measures and Rigorous Discussion of Conditional Kaniadakis Entropy, Int. J. Sci. 4(10): in print.

Amelia Carolina Sparavigna, ORCID 0000-0003-4502-8974

## Information about this Article

This Article was published on 9th November, 2015 at 19:30:37 and has been viewed 3 times.



This work is licensed under a [Creative Commons Attribution 2.5 License](#).

**The full citation for this Article is:**

Sparavigna, A. (2015). Tsallis and Kaniadakis Entropic Measures in Stellar Polytropes. *PHILICA.COM Article number 541*.

[<< Go back](#)

[Review this Article](#)

[Printer-friendly](#)

[Report this Article](#)

Website copyright © 2006-07 Philica; authors retain the rights to their work under [this Creative Commons License](#) and reviews are copyleft under the [GNU free documentation license](#).  
Using this site indicates acceptance of our [Terms and Conditions](#).

This page was generated in 0.2568 seconds.